

# **PREDICTION MODEL OF STUNTING CAUSES IN TODDLERS AGES 12 TO 60 MONTHS IN BAGAN DELI VILLAGE, MEDAN CITY**

**Nenni Dwi APRIANTI LUBIS**

*Universitas Sumatera Utara, Medan, Indonesia*

## **1. Introduction**

The occurrence of short toddlers is one of the nutritional problems experienced by children in the world today. In 2017, around 22.2% or 150.8 million children under five in the world experienced stunting. It was found that more than half of children under five who were stunted in the world came from the Asian continent (55%), and more than one third (39%) were in countries in the African continent. Based on Child Stunting Data Visualizations dashboard prevalence data collected by the World Health Organization (WHO) in 2017, Indonesia is included in the third country after India and Timor Leste with the highest prevalence in the Southeast Asian region. The average incidence of stunting toddlers in Indonesia in the period



2005-2017 reached 36.4% and currently the prevalence of short toddlers in Indonesia tends to be static (Center for Health Data and Information, 2018).

Stunting is a condition where a toddler has less length or height when compared to his age. Nutrient intake in infants becomes very important in supporting growth so that growth faltering does not occur, which can cause stunting. In 2017, 43.2% of children under five in Indonesia experienced an energy deficit, and 28.5% experienced a mild deficiency. For protein adequacy, 31.9% of children under five have a protein deficit and 14.5% have a mild deficit (Directorate of Community Nutrition, 2018).

The socio-economic and educational conditions of parents were considered to be related to the incidence of stunting. The socioeconomic and educational requirements of parents were deemed to be associated with the prevalence of stunting. Based on data from the Joint Child Malnutrition Estimates in 2018, countries with middle to upper income can reduce stunting by up to 64%, whereas in the middle to lower-income countries can only mitigate around 24% from 2000 to 2017. Conversely, in countries with low incomes, there has been an increase in stunting in 2017 (Center for Health Data and Information, 2018). Parental education is a preventative factor for stunting. Highly educated parents have more knowledge about health, nutrition, proper child care, health services, hygiene, adequate food for children, and so on. Higher educated parents also tend to try to increase household income to support the provision of nutritious food in the family (Sultana, Rahman, and Akter, 2019).

Health conditions and nutritional status of mothers before and during pregnancy and after delivery significantly affect fetal growth and risk of stunting. Other factors, such as the mother's short posture, the distance of pregnancy that is too close, the lack of knowledge of the mother, and lack of nutrition during pregnancy can also cause stunting in children. Besides, food obtained by children from birth positively affects their growth, including the risk of stunting. Likewise, the initiation of early breastfeeding, the failure of exclusive breastfeeding, and the process of early weaning can be a factor in stunting. For children who have received complementary foods, it is crucial to consider the quantity and quality of food, food safety, and the diversity of food provided. This study aims to determine the factors associated with the incidence of stunting in infants aged 12 to 60 months in Bagan Deli Village, Medan City.

## 2. Method

This type of research is an observational analytic cross-sectional design. The population in this study were all infants aged 12-60 months who were in Neighborhood VII, Bagan Deli Village, Medan City. Samples were taken in total sampling as many as 59 people. Mothers of toddlers are referred to as respondents.

The independent variables in the study are the mother's education, family income per month, the mother's behavior in serving food, and children's food diversity. Independent variables were measured through interviews using a questionnaire. Meanwhile, the dependent variable is the incidence of stunting obtained from measurements of children's height and compared with age. It is said stunting is if the length or height of a child is more than minus two standard deviations of the median standard of child growth based on WHO. Primary data tabulated frequency. Furthermore, a multivariate regression test was performed to determine the factors associated with the occurrence of stunting.

### *2.1. Mother's behavior in serving food*

We divide into three categories, namely less, sufficient, and well based on 35 items, including knowledge, attitudes, and actions. Poor behavior, if the answer score <56%, sufficient if the answer score is between 56-75%, and good practice, if the answer score  $\geq 76\%$ .

### *2.2. Food diversity of toddlers*

We use the food frequency questionnaire (FFQ) to determine food diversity. Of the 16 food groups categorized in the Guidelines for Measuring Household and Individual Dietary Diversity Score (IHDDS) (FAO, 2010), they are grouped back into nine food groups referring to the Food Diversity Pattern of the Indonesian people. Based on the daily food group consumed by children, we distinguish between the food diversity for less ( $\leq 3$  food groups), moderate (4-6 food groups), and high ( $\geq 7$  food groups).

## 3. Result

Table 1 illustrates the distribution of respondent characteristics. The age range of respondents is  $30.1 \pm 1.4$  years. Based on the mother's last education, a total of 48 respondents completed high school or equivalent (81.4%) and overall worked as a housewife. Most of the family heads work as unskilled laborers (54 people, 91.5%), and five others work as fishermen (8.5%). Family income is mostly in the range of Rp. 1,200,000 to Rp. 2,799,999 (34 respondents, 57.6%). This figure is in the range

of the Medan City Minimum Wage in 2017, which is Rp 2,528,815. The majority of respondents behaved sufficiently while serving food for their children, and only 17 respondents behaved well.

**Table 1.** Characteristics of respondents

Variables		n (%)
Mother's education	Elementary school	4 (6.8)
	Junior school	7 (11.8)
	High school	48 (81.4)
Mother's occupation	Housewife	59 (100)
Husband's occupation	Unskilled laborer	54 (91.5)
	Fishermen	5 (8.5)
Family's income	< Rp. 1,200,000	21 (35.6)
	≥ Rp. 1,200,000-2,799,999	34 (57.6)
	≥ Rp. 2,800,000	4 (6.8)
Mother's behavior in serving food	Poor	14 (23.7)
	Sufficient	28 (47.4)
	<b>Good</b>	<b>17 (28.9)</b>

Most of the respondents' toddlers were female (38 people, 64.4%), with ages between  $24.1 \pm 5.6$  months. The mean weight and height of toddlers were  $11.3 \pm 3.2$  Kg and  $80.3 \pm 21.2$  cm, respectively. Of 59 respondents, the majority of children's nutritional status based on body height index per age is normal, that is 33 toddlers (55.9%), the rest is classified as stunting (26 toddlers, 44.1%). The food diversity of thirty toddlers is classified as moderate. Only 16 toddlers are categorized as high in food diversity because they consume more than seven food groups.

**Table 2.** Characteristics of respondent's children

Variables		n (%)
Gender of toddlers	Male	21 (35.6)
	Female	38 (64.4)
Nutritional status of toddlers (body height per age)	Stunting	26 (44.1)
	Normal	33 (55.9)
Food diversity of toddlers	Less	13 (22)
	Moderate	30 (50.9)
	<b>High</b>	<b>16 (27.1)</b>

From Table 3, it is known that the significant value in the F test is 0.001 so it can be concluded that the mother's behavior in serving food, food diversity, family income, and mother's education together influence the incidence of stunting. Based on the coefficient of determination obtained 0.380 means that the variables  $X_1$ ,  $X_2$ ,  $X_3$ , and  $X_4$  simultaneously affect the Y variable by 38%. The rest is influenced by other variables outside the variables studied.

**Table 3.** Summary of multiple regression analysis

Variable	Regression coefficients	$t_{hitung}$	Sig.
Constant	0.450		
$X_1$ Mother's behavior in serving food	-0.849	- 2.324	<b>0.024</b>
$X_2$ Food diversity of toddlers	1.392	3.886	<b>0.001</b>
$X_3$ Family's income	0.353	1.204	<b>0.234</b>
$X_4$ Mother's education	-0.088	-0.378	<b>0.707</b>
$F_{hitung} = 8.271$			<b>0.001</b>
<b>R Square = 0.380</b>			

Based on multiple regression tests obtained by the regression equation as follows  $Y = 0.450 - 0.8499X_1 + 1.392X_2 + 0.353X_3 - 0.088X_4$ . Toddlers in Bagan Deli Village have stunting opportunities of 0.849 times lower in mothers who behave well in serving food and 0.088 times lower in mothers with high education. Conversely, the risk of stunting will increase 1.392 times in toddlers who consume less variety of food. Likewise, in children under five from low-income families, 0.353 times have a chance of being stunted.

#### 4. Discussion

The actions of parents, especially mothers in consuming healthy foods, have an important role in shaping healthy eating patterns in children. Children's health is something that needs to be taken seriously by parents. For this reason, proper feeding practices should be pursued to keep children healthy. Children's health can be achieved through balanced feeding efforts in accordance with nutritional needs. The role of the mother as the primary caregiver of her child is very necessary starting

from the purchase to the serving of food. If the mother's education and knowledge are low, it is assumed that she is unable to choose to serve food to the family meeting the requirements of balanced nutrition (Rahayu et al, 2015). Parental education, especially for mothers, is a factor associated with the incidence of stunting in infants. This result is consistent with the results of Ni'mah, and Nadhiroh's research (2015) with ( $p = 0.029$ ) and OR of 3.378, which shows that more than half of mothers with stunting under five have low levels of education (61.8%) and more than half of mothers in the group healthy toddlers with high school (67.6%). Research with a sample of toddlers aged 0-2 years in North Hulu Sungai District also shows that low maternal education increases the risk of stunting in children by 1.6 times (Budiastutik, and Rahfiludin, 2019).

Black et al (2008) state that stunting can occur in children over the age of 6 months who do not get enough food both in terms of nutrition and food diversity. Age 6-23 months is a crucial period in stunting prevention. During this time, the child needs enough nutrients to grow and develop. Widyaningsih, Kusnandar, and Anantanyu (2018) showed a significant relationship between food diversity and the incidence of stunting in children under five ( $p < 0.05$ ). As many as 85.4% of children under five with no varied food intake were stunted, and only 14.6% of children under five with food consumption were stunted. When viewed from the distribution of food groups consumed, there is a tendency that stunting toddlers to have a lower intake of green vegetables, vegetables, and fruit sources of vitamin A, vegetables, and other fruits, nuts, and milk when compared to toddlers with normal nutritional status. The percentage of stunting toddlers who consume meat and fish is higher when compared to toddlers with normal nutritional status, but the difference is only 3.2%.

In the Rakotonirainyet et al study (2018) in the Moramanga and Morondava regions, Russia concluded that a mother's education was related to the diversity of food consumed. Children from families with low economic status and mothers without formal education need attention in efforts to feed children. Low socio-economic households are at risk of providing a variety of children's food. The results of Paramashanti, Paratmanitya, and Marsiswati's (2017) research in Bantul Regency, Yogyakarta in children aged 6-24 months showed that the IDDS was strongly associated with the incidence of stunting. Toddlers with low food diversity intake have a 16.67 greater chance of experiencing stunting compared to toddlers with high food diversity intake. Solomon, Aderaw, and Tegegne (2017) explained that mothers' education, income per month, mother's knowledge about food diversity are essential factors in providing minimum food diversity. Consumption of at least four food groups consisting of 1 type of animal food, 1 type of vegetable, 1 type of fruit, and 1 type of staple food. The consumption of these foods is known as the Minimum Dietary Diversity and is also known to be significantly related to the mother's education (Beyene, Worku, and Wassie, 2015).

The majority of families earn between Rp 1,000,000-and 2,999,999 which is 34 respondents. Eleven respondents (32.4%) of them had stunting toddlers, and 23 respondents (67.6%) had normal nutritional status. The analysis shows that family income is a factor associated with the incidence of stunting. Ni'mah, and Nadhiroh research (2015) explains that family income is a factor related to the prevalence of stunting ( $p = 0.044$ ) with an OR of 3,250. Family income is one of the socio-economic indicators of the family so that the family can meet the nutritional consumption needs. Budiastutik, and Rahfiludin (2019) state that work and parental income have an impact on the incidence of stunting.

## 5. Conclusion

Mother's education, mother's knowledge, food diversity, and family income are factors that influence the incidence of stunting in infants. Therefore it is necessary to have a program that supports overcoming the prevalence of stunting in infants from both the government and non-government sectors, namely by improving nutrition education and counseling to mothers and regularly monitoring the implementation of household-level balanced nutrition.

## Acknowledgments

Thank you to Dessy Andrianty and Geetha Palanivelu for helping with data collection.

## Disclosure statement

No potential conflict of interest was reported by the author.

## Contact

**E-mail:** nenni@usu.ac.id

## References and notes:

- Beyene, M., Worku, A.G. & Wassie, M.M. “Dietary Diversity, Meal Frequency and Associated Factors among Infant and Young Children in Northwest Ethiopia: A Cross-sectional Study”. *BMC Public Health*, vol. 15, 2015,1007,https://doi.org/10.1186/s12889-015-2333-x
- Black R.E., Allen L.H., Bhutta Z.A., Caulfield, L.E., deOnis, M., Ezzati, M. et al. “Maternal and Child Undernutrition: Global and Regional Exposures and Health Consequences”. *Lancet*, vol. 371, no. 9608, 2008, pp. 243–60,https://doi.org/10.1016/S0140-6736(07)61690-0
- Budiastutik, I. & Rahfiludin, M.Z. “Risk Factors of Child Stunting in Developing Countries”. *Amerta Nutr*, vol.3, no. 3, 2019, pp. 122-126, http://dx.doi.org/10.20473/amnt.v3i3.2019.122-129
- Center for Health Data and Information. 2018. “Stunting in Indonesia”. *Health Data and Information Bulletin*. ISSN: 2088-270X. Ministry of Health of the Republic of Indonesia. Available form:https://www.google.com/url?sa=i&url=https%3A%2F%2Fpusdatin.kemkes.go.id%2Ffolder%2Fview%2F01%2Fstructure-publikasi-pusdatin-buletin.html&psig=AOvVaw0XaYtO0vRvnPnlj95OzS0a&ust=1584780124201000&source=images&cd=vfe&ved=0CA0QjhxqFwoTCLCn8ZvUqOgCFQAAAAAdAAA AABAD
- Directorate of Community Nutrition. 2018. “Nutrition Status Monitoring Results in 2017”. *Nutrition Status Monitoring Handbook for 2017*. Directorate General of Community Health, Ministry of Health of the Republic of Indonesia. Available form: https://www.google.com/url?sa=i&url=http%3A%2F%2Fsehatnegeriku.kemkes.go.id%2Fbaca%2Fnumum%2F20180125%2F3424539%2Fbuku-saku-hasil-pemantauan-status-gizi-psg-tahun2017%2F&psig=AOvVaw2ki2DZyNzoumBnPuX3xcqH&ust=1584780333292000&source=images&cd=vfe&ved=0CA0QjhxqFwoTCNDrgILVqOgCFQAAAAAdAAA AABAD
- FAO. 2010. “Guidelines for Measuring Household and Individual Dietary Diversity”. Available form: https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved=2ahUKEwirqefz8KjoAhUclbcAHTgUDYQQFjABegQIAXAB&url=http%3A%2F%2Fwww.fao.org%2F3%2Fa-i1983e.pdf&usg=AOvVaw3yikMCPfWcrH-3OvYKakT5
- Ni'mah, K. & Nadhiroh, S.R. 2015. “Factors Related to Stunting on Toddlers”. *Media Gizi Indonesia*, vol. 10, no.1, 2015, pp. 13-19, http://dx.doi.org/10.20473/mgi.v10i1.13-19
- Paramashanti, B.A., Paratmanitya, Y. & Marsiswati, M. “Individual Dietary Diversity is Strongly Associated with Stunting in Infants and Young Children”. *The Indonesian Journal of Clinical Nutrition*, vol. 14, no. 1, 2017, 19-26,https://doi.org/10.22146/ijcn.15989
- Rahayu, A., Yulidasari, F., Putri, A.O., Rahman, F. “Birth Weight Records with Stunting Incidence among Children under Two Years Old”. *National Public Health Journal*, vol. 10, no. 2, 2015, pp. 67-73, http://dx.doi.org/10.21109/kesmas.v10i2.882
- Rakotonirainy, N.H., Razafindratovo, V., Remonja, C.R., Rasoloarijaona, R., Piola, P., Raharintsoa, C. & Randremanana, R.V. “Dietary Diversity of 6- to 59-Month-Old Children in Rural Areas of Moramanga and Morondava Districts, Madagascar”. *PLoS One*, vol.13, no.7, 2018, e0200235, doi: 10.1371/journal.pone.0200235
- Solomon, D., Aderaw, Z. & Tegegne, T.K. “Minimum Dietary Diversity and Associated Factors among Children Aged 6–23 Months in Addis Ababa, Ethiopia”. *Int J Equity Health*, vol. 16, no.181, 2017, https://doi.org/10.1186/s12939-017-0680-1
- Subasi, A. Subasi.G. (2021). Innovation in health management and social care. *International Journal of Humanities and Social Development Research*. Volume 5, Number 2. pp. 7-19. https://doi.org/10.30546/2523-4331.2021.5.2.7
- Sultana, P., Rahman, M.M. & Akter, J. “Correlates of Stunting among Under-Five Children in Bangladesh: A Multilevel Approach”. *BMC Nutr*, vol. 5, no. 41, 2019, https://doi.org/10.1186/s40795-019-0304-9
- Widyaningasih, N.N., Kusnandar, Anantanyu, S. “Food Diversity, Parenting Practices and Stunting in Toddlers Age 24-59 Months”. *The Indonesian Journal of Nutrition*, vol. 7, no. 1, 2018, pp. 22-29, https://doi.org/10.14710/jgi.7.1.22-29

**Cite this article as:** APRIANTI LUBIS Dwi Nenni. Prediction model of stunting causes in Toddlers ages 12 to 60 months in Bagan Deli village, Medan city. *Journal of Science and Innovative Technologies*. Number 21, 2022. pp..55-65. DOI: 10.30546/2616-4418.21.2022.55



## XÜLASƏ

### **Medan şəhərinin Baqan Deli qəsəbəsində 12-60 aylıq körpələrdə böyümənin geriləmə səbəblərinin proqnozlaşdırılması modeli**

**Nenni Dwi APRIANTI LUBİS**

*Sumatera Utara Universiteti Medan, İndoneziya*

Körpələrin önəmli dövründəki qida çatışmazlığı, sonrakı dövrdə qida ehtiyacları qarşılansa belə, qalıcı problem yarada bilər. Qida çatışmazlığı hallarının yüksək olmasında anaların qidalanma haqqında məlumatının aşağı olması, ananın müəyyən qidalara olan inamı və ailə gəlirinin aşağı olması kimi səbəblər göstərilir. Bu araşdırma 12-59 aylıq körpələrdə boy artımının azalması hallarına təsir edən amilləri müəyyən etmək məqsədi daşıyan analitik -tədqiqatdır.

Bu tədqiqatın əsas araşdırma istiqamətləri Medan şəhərinin Baqan Deli qəsəbəsinin əhalisi və VII məhəllədə yaşayan 12-60 aylıq körpələr idi. Beş yaşından kiçik uşaqları olan analar bu tədqiqatın respondentlərdir və ümumi seçmə zamanı bütün əhali arasından 59 nəfər seçilmişdir. Ananın qida təminatında və uşaqların qida müxtəlifliyi ilə bağlı araşdırma məlumatları beş yaşından kiçik uşaq anaları ilə məlumatlar müsahibə şəklində aparılaraq toplanmışdır. Bundan əlavə, beş yaşdan kiçik uşaqların qidalanma vəziyyəti boy ölçüləri ilə əldə edilir.

Alınan məlumatlar müəyyən standartlara əsaslanaraq, kateqoriyalara uyğun olaraq işlənmiş və təhlil edilmişdir. Nəticələr göstərdi ki, respondentlərin orta yaş həddi  $30,1 \pm 1,4$  ildir və onların əksəriyyəti orta məktəbi bitirmişdir və ailənin gəliri ümumilikdə ayda təxminən-1.200.000-2.799.999.Rp arasında dəyişir.

Uşaqlara yemək verməkdə respondentlərin yalnız 28,9%-i müəyyən qədər yaxşı nəticə göstərmiş, qalanları yəni, 23,7% və 47,4%-i isə yaxşı nəticə göstərmədilər. Ümumiyyətlə, respondentlər uşaq yeməklərinin təqdim edilməsində olduqca fərqlidirlər (50,9%) .

Yaş indeksinə görə uşaqların qidalanma vəziyyətinə əsasən, uşaqların 44,1%-i inkişafdan geri qalmış, qalanları isə normaldır. Çoxsaylı reqressiya analizinin nəticələri  $Y=0,450-0,849X_1+1,392X_2+0,353X_3-0,008X_4$  tənliyini 0,001 əhəmiyyətlik dəyəri ilə göstərilmişdir. Odur ki, belə nəticəyə gəlmək olar ki, ananın yemək təminatında davranışı, uşağın qida müxtəlifliyi, ailənin gəliri, ananın təhsili böyümənin geriləmə hallarına çox təsir edir.

**Açar sözlər:** Ailə gəliri, qida müxtəlifliyi, ananın davranışı, ananın təhsili, boy artımı

## РЕЗЮМЕ

### **Модель для прогнозирования причин задержки роста у детей в возрасте 12–60 месяцев в Баган Дели, Медан**

**Ненни Дви АПРИАНТИ ЛУБИС**

*Университет Суматера Утара, Медан, Индонезия*

Недоедание у младенцев в критические периоды может создать остаточные проблемы, даже если потребности в питании будут удовлетворены в более позднем периоде. Высокий уровень недоедания объясняется низкой осведомленностью матерей о питании, низким доверием матерей к определенным продуктам питания и низким доходом семьи.

Настоящее исследование представляет собой аналитическое исследование, направленное на выявление факторов, влияющих на снижение увеличения роста у детей в возрасте 12–59 месяцев. Основными объектами исследования в данной статье были жители поселка Баган-Дели в городе Медан и младенцы в возрасте 12-60 месяцев, проживающие в 7-м районе. Респондентами в данном исследовании были матери с детьми в возрасте до пяти лет, а в ходе отбора среди общего населения было отобрано 59 человек. Данные исследований питания матерей и разнообразия питания детей были собраны в форме интервью с матерями детей до пяти лет. Кроме того, состояние питания детей в возрасте до пяти лет определяется размером роста. Полученные данные были обработаны и проанализированы по категориям на основе определенных стандартов. Результаты показали, что средний возраст респондентов составил  $30,1 \pm 1,4$  года, большинство из них окончили среднюю школу, а доход семьи в целом колебался примерно в пределах 1 200 000-2 799 999 рупий в месяц. Только 28,9% респондентов показали хорошие результаты в кормлении детей, а остальные, т.е. 23,7% и 47,4%, не показали хороших результатов.

В целом респонденты сильно различаются по предоставлению детского питания (50,9%). Основываясь на состоянии питания детей по возрастному показателю, 44,1% детей отстают в развитии, а остальные развиваются в норме.

Результаты множественного регрессионного анализа показывают уравнение  $Y = 0,450 - 0,849X_1 + 1,392X_2 + 0,353X_3 - 0,008X_4$  со значением значимости 0,001.

Следовательно, можно сделать вывод, что существенное влияние на задержку роста оказывают поведение матери в обеспечении пищей, разнообразие питания ребенка, доход семьи, образование матери.

**Ключевые слова:** Доход семьи, разнообразие продуктов питания, поведение матери, образование матери, рост, рост